

*Citation for published version:*

Longrich, N, Pereda-Suberbiola, X, Pyron, RA & Jalil, NE 2021, 'The first duckbill dinosaur (Hadrosauridae: Lambeosaurinae) from Africa and the role of oceanic dispersal in dinosaur biogeography', *Cretaceous Research*, vol. 120, 104678. <https://doi.org/10.1016/j.cretres.2020.104678>

*DOI:*

[10.1016/j.cretres.2020.104678](https://doi.org/10.1016/j.cretres.2020.104678)

*Publication date:*

2021

*Document Version*

Peer reviewed version

[Link to publication](#)

*Publisher Rights*

CC BY-NC-ND

**University of Bath**

**Alternative formats**

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Journal Pre-proof

The first predatory dance fly of the subfamily Ocydromiinae with specialized, raptorial legs in mid-Cretaceous amber from Myanmar (Diptera: Hybotidae)

Valerie Ngô-Muller, Michael S. Engel, Romain Garrouste, Jean-Marc Pouillon, André Nel



PII: S0195-6671(20)30384-0

DOI: <https://doi.org/10.1016/j.cretres.2020.104697>

Reference: YCRES 104697

To appear in: *Cretaceous Research*

Received Date: 6 May 2020

Revised Date: 7 September 2020

Accepted Date: 2 November 2020

Please cite this article as: Ngô-Muller, V., Engel, M.S., Garrouste, R., Pouillon, J.-M., Nel, A., The first predatory dance fly of the subfamily Ocydromiinae with specialized, raptorial legs in mid-Cretaceous amber from Myanmar (Diptera: Hybotidae), *Cretaceous Research*, <https://doi.org/10.1016/j.cretres.2020.104697>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Elsevier Ltd. All rights reserved.

Ngô-Muller wrote the paper, prepared the specimen, and made the photos

Engel wrote the paper

Garrouste wrote the paper

Pouillon wrote the paper and prepared the specimen

Nel wrote the paper, searched the references, and made the analyses

**The first predatory dance fly of the subfamily Ocydromiinae with specialized, raptorial legs in mid-Cretaceous amber from Myanmar (Diptera: Hybotidae)**

Running head

mid-Cretaceous hunter dance fly

Valerie Ngô-Muller<sup>a,b</sup>, Michael S. Engel<sup>c</sup>, Romain Garrouste<sup>b</sup>, Jean-Marc Pouillon<sup>d</sup>, André Nel<sup>b,\*</sup>

<sup>a</sup>*UFR Sciences du Vivant, Université Paris Diderot, Université de Paris, Paris, France*

<sup>b</sup>*Institut de Systématique, Évolution, Biodiversité, ISYEB - UMR 7205 – CNRS, MNHN, UPMC, EPHE, Muséum national d'Histoire naturelle, Sorbonne Universités, Université des Antilles, 57 rue Cuvier, CP 50, Entomologie, F-75005, Paris, France*

<sup>c</sup>*Division of Entomology, Natural History Museum, and Department of Ecology & Evolutionary Biology, 1501 Crestline Drive – Suite 140, University of Kansas, Lawrence, Kansas 66045, USA*

<sup>d</sup>*Nivolas Vermelle, France*

\* Corresponding author. E-mail: [anel@mnhn.fr](mailto:anel@mnhn.fr)

LsidXXX (provided before manuscript acceptance)

**ABSTRACT**

The first ocydromiine hybotid fly is described and illustrated from a remarkable male preserved in mid-Cretaceous amber from northern Myanmar. *Pouillonhybos venator*, gen. et

sp. nov., is distinguished from other members of the subfamily Ocydromiinae as well as other lineages of living and fossil Hybotidae. The holotype of *P. venator* exhibits spectacular specializations of the mid and hind legs, modifications likely linked to the grasping of prey either during capture and/or while feeding. The species reported here is the earliest evidence of significant leg modifications in Hybotidae indicating an early appearance of such specializations in the family's history.

**Keywords:** Brachycera; Cenomanian; Empidoidea; predatory dance flies; taxonomy

## 1. Introduction

The family Hybotidae largely includes predaceous flies that typically frequent on fallen logs or semi-aquatic vegetation in forested environments. Although generally predators, a few are known to feed on pollen. Where known for the predatory taxa, many species capture preys while in flight although tachydromiines more readily subdue their prey on plant surfaces (Chvála, 1983). During feeding, hybotids grasp the prey item with their hind legs while manipulating their food and stabilizing themselves with their fore and midlegs (Wilder, 1974). Accordingly, various modifications of the hind legs have arisen among Hybotidae which appear to be associated with capturing and holding their prey, the most elaborate of which belong to groups believed to take their victims while in flight. The fossil record of the family is scanty, with comparatively few records (Evenhuis, 1994). Hybotid flies are quite rare in the Mesozoic, with only the Early Cretaceous genus *Pseudoacarterus* Waters, 1989, the mid-Cretaceous *Electrocyrtona* Cockerell, 1917, and the Late Cretaceous *Archiplatypus* Kovalev, 1974. Herein we report from the mid-Cretaceous Burmese amber a remarkable new hybotid fly of the subfamily Ocydromiinae with highly specialized male hind legs, implying that the

evolution of such features appeared early in the family's history. We provide a description of this new taxon and comparisons with the diversity of modern and other extinct genera.

## 2. Material and methods

The specimen comes from the earliest Cenomanian amber ( $98.79 \pm 0.62$  Ma, based on U-Pb zircon dating of the volcanoclastic matrix; Shi et al., 2012); location map of amber-bearing locality in Yin et al. (2018: fig. 1A). It is preserved in a piece of clear, yellow amber. The amber piece was cut, shaped, and polished using a diamond disk under water and diatomite, and was then mounted between two coverslips with sugar medium dissolved at saturation in water, before being examined and photographed. The inclusion was examined and measured under incident light with an Olympus SZX9 and Leitz Wetzlar binocular microscopes. The holotype is deposited in the amber collection of the Musée d'Histoire Naturelle et d'Ethnographie de Colmar (MHNEC), France under collection number MHNE.2020.7.1. We follow the terminology of Cumming & Wood (2009) and the empidoid classification of Sinclair & Cumming (2006).

## 3. Systematic palaeontology

Order Diptera Linnaeus, 1758

Superfamily Empidoidea Latreille, 1809

Family Hybotidae Meigen, 1820

Subfamily Ocydromiinae Schiner, 1862

Genus *Pouillonhybos* Ngô-Muller, Engel & Nel, gen. nov.

LSIDurn:lsid:zoobank xxxx

Type species: *Pouillonhybos venator* Ngô-Muller, Engel & Nel, sp. nov.

*Diagnosis.* Proboscis short, oriented ventrally. Antenna with stylus elongate, arista-like, longer than postpedicel. Protibial gland present. Mesofemur crassate, with four stout, elongate, ventral setae; mesotibia elongate, basally slightly geniculate, with two outer rows of short, oblique setae and a single, elongate, apical spur. Hind legs greatly modified; metacoxa with one elongate seta; metafemur with pronounced inner curvature, with two stout, thickened, elongate subapical spines and a series of stout, elongate setae ventrally as well as a row of shorter, curved prolateral (ectal) setae; metatibia strongly arched with a prominent, thick, subapical, hook-like lobe, and a row of long inner setae, and an apical spur; metabasitarsus broadened, with an ectal row of curved setae and an ental row of long, rather straight setae. Wings well developed, much longer than abdomen; pterostigma lacking; C terminating at wing apex, on  $M_1$ ; stem of Rs short;  $R_{2+3}$  comparatively straight;  $R_{4+5}$  and  $M_1$  roughly parallel; *dm* present, elongate; m-m arched, with faint nebulous trace of  $M_2$ , trace disappearing after length about  $0.5 \times$  m-m length; two veins emitted from *dm* reaching wing margin ( $M_1$  and  $M_4$ ); *cua* present, elongate, about as long as *bm*, with apex oblique, with outer angle obtuse, without spur vein; vein dm-m strongly oblique; CuA not broadly arching apically to CuA+CuP; CuP and CuA+CuP faint, not reaching wing margin; anal lobe narrow; alula absent. Epandrium with pair of articulated surstyli apically.

*Etymology.* The new generic name is a combination of the surname Pouillon, honoring Jean-Marc Pouillon, who donated the type specimen, and *Hybos* Meigen, the type genus of the family. The gender of the name is masculine.

***Pouillonhybos venator*** Ngô-Muller, Engel & Nel, sp. nov.

Figs 1–3

LSIDurn:lsid:zoobank xxxx

*Etymology.* The specific epithet is taken from the Latin word '*venator*', meaning, 'hunter', referring to the specialized metatibiae of this species.

*Material.* Holotype MHNE.2020.7.1 (a complete specimen in amber, coll. Jean-Marc Pouillon); Musée d'Histoire Naturelle et d'Ethnographie de Colmar, France.

*Horizon and locality.* Noiye Bum Hill, Hukawng Valley, Kachin State, Myanmar; lower Cenomanian, mid-Cretaceous.

*Diagnosis.* As for genus; only one included species.

*Description.* Male, body 2.03 mm long (Fig. 1); head 0.22 mm long, 0.30 mm wide; compound eyes apparently setose, with few visible very short setae, large, meeting at one point above (a gas bubble occludes views of the upper part of the head); ptilinal fissure absent; occiput and frons bare; apex of antenna with lengthened, arista-like stylus (Fig. 3A), possibly dimerous as a very short structure is present between it and postpedicel, apical on postpedicel; postpedicel pear-shaped, 0.06 mm long, quite shorter than stylus (0.66 mm long) and shorter than pedicel; pedicel with an apical crown of strong setae, longer than scape; proboscis short, oriented ventrad.

Thorax 0.64 mm long, 0.45 mm wide, with bare laterotergite; two elongate setae on scutellum and three near base of wing.

Wing hyaline (Fig. 1B), longer than abdomen, apex rounded, 2.22 mm long, 0.45 mm wide, covered with microtrichia, without darkened pterostigma; Sc ending on C, 1.08 mm from wing base; C extending to wing apex, terminating at apex of M1; Rs originating well distad level of humeral vein (h), 0.39 mm from it; branches of Rs not strongly thickened and crowded anterobasally; R2+3 elongate and straight, 0.08 mm long; R4+5 unbranched, 1.29 mm long, nearly straight and parallel to M1; cell *dm* present, 0.69 mm long, 0.13 mm wide, with three veins distally emerging from it: M1 (0.6 mm long), M2 (present as an exceedingly faint nebulous vein disappearing well before wing margin, distinctly shorter than m-m



crossvein), and M4 (0.19 mm long); CuA forming a distinct angle with basal part of CuP; cell *cua* scarcely shorter than cell *bm* (*bm-cu* and CuA nearly confluent), with outer angle obtuse; CuA+CuP vein not reaching wing margin; narrow anal lobe, 0.13 mm wide; lunule absent; a series of long setae on anal margin; haltere elongate, 0.41 mm long, 0.08 mm wide.

Legs (Figs 2, 3A). Pretarsal claws simple, empodia setiform; protibial gland present; profemur 0.29 mm long, 0.06 mm wide, curved, with a series of at least five long setae along inner surface; protibia 0.29 mm long, 0.04 mm wide, with three rows of stout setae, one apical spur; probasitarsus elongate, 0.18 mm long, with two series of long setae; protarsomere II 0.11 mm long; protarsomere III 0.08 mm long; protarsomere IV 0.04 mm long; protarsomere V 0.06 mm long; mesofemur crassate, approximately 0.42 mm long, 0.06 mm wide, with four stout, thickened, erect, spine-like setae on ventral surface; mesotibia 0.53 mm long, 0.04 mm wide, basally geniculate, with two series of rather short, oblique setae, a single elongate apical spur; mesotarsus ca. 0.53 mm long; metacoxa with a single, elongate, posteroventral seta, 0.29 mm long; metafemur 0.76 mm long, 0.09 mm wide, with a pronounced inner curvature and two stout, thickened, enlarged, subapical spurs (0.16 mm long) and a series of long, stout, erect, spine-like setae along ventral surface, a series of shorter, finer, curved, oblique setae along dorsal surface (Fig. 2); metatibia basally geniculate, strongly curved, with a subapical hook-like lobe (0.1 mm long), apex of lobe narrowly rounded, a row of long, oblique setae on inner surface, a single, short, basally thickened apical spur, outer surface with a row of erect to semierect, short, fine setae; metabasitarsus greatly broadened and somewhat trapezoidal as lateral margins slightly converge apically, 0.09 mm long, 0.04 mm wide, outer edge with a row of short, strongly curved setae, inner edge with a row of longer, obliquely erect, largely straight setae; metarsomere II elongate, 0.16 mm long, 0.01 mm wide, with two rows of fine, long setae; metatarsomere III 0.09 mm long, 0.01 mm wide, metatarsomere IV 0.08 mm long, 0.008 mm wide, metatarsomere V 0.08 mm long, 0.008 mm wide.

Abdomen elongate (Fig. 2), rather narrow, 1.0 mm long, 0.24 mm wide, with a single, elongate spine-like seta laterally on segments II–V; male terminalia asymmetrical and rotated 90° (Fig. 3B); cercus thin with a small inner extension; hypoproct small near cercal base; phallus large, as long as and broader than cerci; epandrium greatly enlarged, spoon-shaped, apparently with an apical pair of articulated surstyli.

Female unknown.

#### 4. Discussion

*Pouillonhybos* gen. nov. can be placed within the Empidoidea owing to the following characters: ptilinal fissure and lunule absent; empodia setiform; pedicel shorter than flagellum; flagellum with three flagellomeres; cell *cua* small; stylus elongate and thin; wing apex rounded; branches of R not strongly thickened and crowded anterobasally; C extending to wing apex; CuA+CuP not reaching wing margin. Using those traits outlined by recent authors (*e.g.*, Steyskal & Knutson, 1981; Sinclair & Cumming, 2006, 2017), the new fossil genus falls among the Hybotidae based on the following characters: apex of antenna with lengthened, bristle-like stylus (stylus also not setulose); thorax with bare laterotergite; protibial gland present; C ending at wing apex, instead of circumambient (but some empidids also have a costa terminating at the wing apex); Rs originating well distal to level of humeral vein (h); R4+5 unbranched; CuA forming a distinct angle with basal part of CuP. It should be noted that some Empididae have a wing venation quite similar to that of *Pouillonhybos* (*viz.* *Macrostomus* Wiedemann, 1817), especially in the narrow anal lobe and shape of cell *cua*, and even in the C terminating on M1. Such empidids, however, differ from *Pouillonhybos* in the setulose scape (not so in the fossil), shape of the more elongate flagellum, and, in the case of *Macrostomus*, lacking a defined stylus (Smith, 1961: fig. 3; Rafael & Cumming, 2004: figs 1, 3).

Wahlberg & Johanson (2018) presented a molecular phylogenetic analysis of the Empidoidea, in which Hybotidae are subdivided into six subfamilies: Bicellariinae Sinclair & Cumming, 2006; Ocydromiinae Schiner, 1862; Oedaleinae Chvala, 1983; Hybotinae Meigen, 1820; Tachydromiinae Meigen, 1822; and Trichininae Chvala, 1983. Most recently, Sinclair (2019) added the subfamily Stuckenbergomyiinae Sinclair, 2019. Affinities between the new fossil and Trichininae (*Trichinomyia* Tuomikoski, 1959 and *Trichina* Meigen, 1830) are excluded because of the absence of an alula, the narrow anal lobe, the absence of a pterostigma, and the elongate stylus (Tuomikoski, 1959). The Tachydromiinae are excluded because  $M_2$  is present, albeit only as a short, faintly nebulous vein, cell *dm* present, and cell *cua* elongate (Sinclair & Cumming, 2006, 2017). Similarly, the subfamily Bicellariinae (*Bicellaria* Macquart, 1823) can be discounted owing to the narrow anal area and presence of cell *dm* (Kato, 1971). The Oedaleinae have a branched  $M_{1+2}$  and an anterad projecting proboscis, quite unlike *Pouillonhybos*, and the Stuckenbergomyiinae differ from the new fossil genus in the truncate cell *cua*, shortened stylus, and three veins emitted from cell *dm*.

Affinities with the Hybotinae would also be excluded because of the presence of  $M_2$  in *Pouillonhybos*, but as it is rudimentary in the present fossil we prefer to compare the new genus to extant and fossil hybotine genera. The short proboscis of *Pouillonhybos* excludes affinities with *Syneches* Walker, 1852 (incl. *Parahybos* Kertesz, 1899, *Harpamerus* Bigot, 1859, and *Epiceia* Walker, 1860), *Hybos* Meigen, 1803, *Smithybos* Ale-Rocha, 2000, *Syndyas* Loew, 1857, and *Lactistomyia* Melander, 1902. In the genera *Afrohybos* Smith, 1967, *Cerathybos* Bezzi, 1909, *Euhybus* Coquillett, 1895, and *Neohybos* Ale-Rocha & Carvalho, 2003 cell *cua* extends far distal to cell *bm* (Melander, 1902; Smith, 1967; Ale-Rocha, 2000, 2002, 2008; Ale-Rocha & Carvalho, 2003; Grootaert & Yang, 2009; Sinclair, 1996, 2011; Liu et al., 2012, 2014), quite unlike *Pouillonhybos* in which these cells are about subequal in length, with *cua* only scarcely shorter than *bm*. *Acarterus* Loew, 1858, *Lamachella* Melander,

1928, and the Cretaceous genus *Pseudoacarterus* Waters, 1989 have cell *cua* apically convex (*i.e.*, CuA is convexly arched), while in the new genus CuA is straight and oblique (Waters, 1989; Sinclair, 1996). Additionally, in *Lamachella* *cua* is distinctly shorter than *bm*, a pterostigma is usually present, m-cu is more elongate and not in near alignment with CuA, and R<sub>2+3</sub> is sinuate, unlike *Pouillonhybos* (Smith, 1969: figs 137-138; Sinclair & Cumming, 2017). *Chillcottomyia* Saigusa, 1986, *Stenoproctus* Loew, 1858, and the Miocene genus *Syneproctus* Solórzano-Kraemer et al., 2020 have cell *cua* quite shorter than *bm*, and with apex convex, nearly perpendicular to the two veins CuA and CuP, unlike *Pouillonhybos* (Smith, 1969: figs 127-128, 130; Yang & Grootaert, 2004; Solórzano-Kraemer et al., 2020).

The fossil shares many traits with the subfamily Ocydromiinae, in which cell *cua* is shorter than or about as long as cell *bm*, with its outer angle obtuse; the postpedicel shorter than the arista-like stylus; a cell *dm* present; and the proboscis oriented ventrad, all as in *Pouillonhybos* (Sinclair & Cumming, 2000, 2006; Wahlberg & Johanson, 2018). In addition, ocydromiines also have an epandrium with an apical pair of articulated surstyli, another trait apparently present in *Pouillonhybos*, pointing to a placement within the subfamily for this new Cretaceous genus.

Within this subfamily, *Leptopezella* Sinclair & Cumming, 2007 has no discal cell, a pterostigma, cell *cua* clearly shorter than *bm*, and no metatibial extension, and is therefore quite different from *Pouillonhybos* (Sinclair & Cumming, 2007). *Abocciputa* Plant, 1989 and *Chvalaea* Papp & Földvári, 2001 differ from *Pouillonhybos* in that cell *cua* is significantly shorter than cell *bm*, there is no trace of M<sub>2</sub>, and R<sub>4+5</sub> is not parallel to M<sub>1</sub> (Plant, 1989; Papp & Földvári, 2002; Shamshev et al., 2017), and the genera *Austropeza* Plant, 1989, *Pseudoscelolabes* Collin, 1933, and *Leptodromiella* Tuomikoski, 1936 also have cell *bm* much longer than cell *cua* (Tuomikoski, 1936; Collin, 1926, 1928, 1933; Chvála, 1983; Plant, 1989). *Oropezeella* also has a rather long cell *dm* and quite short veins emerging from it, unlike

*Pouillonhybos* (Chvála, 1983; Ale Rocha, 2007). In the genus *Stylocydromia* Saigusa, 1986 there is a long cell *dm*, while the base of *Rs* is much closer to *h* than to *r-m*, unlike the condition in *Pouillonhybos* (Saigusa, 1986). The genus *Leptopeza* Macquart, 1827 has a rudimentary *M*<sub>1</sub> and cell *bm* much longer than cell *cua* (Chvála, 1983), while in *Ocydromia* Meigen, 1820, the stylus is supra-apical rather than apical in *Pouillonhybos*, and moreover cell *bm* is much longer than cell *cua* and there is no trace of a third vein emerging from *dm* (Chvála, 1983). *Apterodromia* Oldroyd, 1949 shares with *Pouillonhybos* a cell *cua* nearly as long as *bm*, but it has an anal area much more reduced than *Pouillonhybos* and a well-defined *M*<sub>2</sub> (Sinclair & Cumming, 2000). *Neotrichina* Sinclair & Cumming, 2000 has the stylus scarcely as long as the postpedicel, three veins emitted from *dm* that reach the wing margin, cell *cua* much shorter than *bm*, an anal vein almost complete to the wing margin, *CuA* closing *cua* almost straight, all quite distinct from the character states present in *Pouillonhybos*; also some species of *Neotrichina* have the metatibia dilated apically but without a lateral expansion, and therefore unlike that in *Pouillonhybos* (Collin, 1933; Sinclair & Cumming, 2000). *Hoplopeza* Bezzi, 1909 and *Scelolabes* Philippi, 1865 can be excluded from having any affinity with the new fossil genus because they have *Rs* short, arising near the apex of cell *bm*, and cell *cua* much shorter than *bm* (Collin, 1933; Rafael, 1995; Sinclair & Cumming, 2000). Lastly, *Leptodromia* Sinclair & Cumming, 2000 has cell *cua* two-thirds length of cell *bm* and a short *M*<sub>1</sub> that does not reach the wing margin (Bezzi, 1904; Sinclair & Cumming, 2000).

Among known extinct genera, the Eocene Baltic amber genus *Palaeoleptopeza* Meunier, 1908, was considered as near to “*Leptopeza* but with spinose hind femora” by Melander (1928: 371), and differs from *Pouillonhybos* in that the postpedicel is as long as the stylus, *R*<sub>2+3</sub> is distinctly arcuate, and *M*<sub>2</sub> is more clearly defined as a fuscous nebulous trace (Meunier, 1908: pl. 7, fig. 15, pl. 8, fig. 4). *Eommocydromia difficilis* Schlüter, 1978 is the

unique Cretaceous taxon currently attributed to the Ocydromiinae; it is based on an incomplete specimen in amber with numerous characters of wing venation missing (in particular, the relative positions of r-m, base of Rs, and h are unknown), rendering it impossible to determine if this assignment is valid or whether it might belong to the Dolichopodidae. It has three long veins emerging from an elongate cell *dm*, which is quite unlike any other Ocydromiinae. The cells *bm* and *cua* are not preserved, and we believe that attribution to Ocydromiinae is difficult to justify (Schlüter, 1978: figs 86-89). *Ecommocydromia difficilis* is perhaps best considered as incertae sedis until more completely preserved specimens are recovered. The Eocene Baltic amber genus *Meghyperella* Meunier, 1908 was considered by Melander (1928: 370) as “apparently related to *Leptopeza*” (Ocydromiinae), and possibly a “predecessor of *Stenoproctus*” (Hybotinae). It differs from *Pouillonhybos* in the exceptionally short stylus (Meunier, 1908), and needs to be revised. Evenhuis (1994) listed it among the fossil ‘Empididae’. Note that the alleged Oligocene hybotine genus *Eternia* Martins-Neto et al., 1992 is based on a poorly preserved fossil, but it preserves *cua* as long as *bm* with an obtuse outer angle, suggesting it belongs to the Ocydromiinae (Martins-Neto et al., 1992). Lastly, the putative Oligocene hybotine genus *Tremembella* Martins-Neto et al., 1992 is based on an exceptionally poorly preserved specimen and until more complete material is discovered it remains impossible to compare with the other genera.

The only other Hybotidae from Burmese amber is *Electrocyrtona burmanica* Cockerell, 1917, a genus that differs from *Pouillonhybos* in the absence of cell *dm*, a much broader cell *bm*, and a cell *cua* much shorter than *bm* (Cockerell, 1917: fig. 5). Melander (1928: 368) indicated that *Electrocyrtona* resembled *Bicellaria*, while Grimaldi & Cumming (1999: 51) considered it as belonging to Tachydromiinae.

## 5. Conclusions

*Pouillonhybos* is the first ocydromiine from Burmese amber and probably the only definitive representative of this subfamily from the Cretaceous. Modern Ocydromiinae are predators (Chvála, 1976), and like other hybotids often hold their prey with the hind legs while feeding. However, no living ocydromiines have a massive prominence on the metatibia like the one present in *Pouillonhybos*. This structure, together with the stout setae and spines on the metafemur, erect and spine-like setae on ventral surface of mesofemur, and setae on the other femora and tibiae, coxae, and abdomen, strongly suggest that *Pouillonhybos* was a predator who, like surviving relatives, captured and hold prey with its hind legs. The hybotine genera *Hybos*, *Syndyas*, and *Syneches* have raptorial hind legs (Grootaert, 1996), with strong spines on the metafemur but no prominent extension on the metatibia. Some species of *Lactistomyia* also have raptorial hind legs with a metatibial lateral extension, albeit situated in a distinctly more proximal position than that of *Pouillonhybos* (Ale-Rocha, 2008). *Afrohybos* has also a highly modified hind leg with a ventral extension of the metafemur and strong spines on the metatibia (Smith, 1969: fig. 96), while the hybotine genus *Syneproctus* has a metatibia with an apical extension most analogous to that of the fossil, but quite a bit smaller than that of *Pouillonhybos*.

Many Hybotidae and Dolichopodidae have raptorial fore and/or mid legs with greatly modified femora and/or tibiae, and some dolichopodids have a strong spine on an apical extension of the meso- and/or metatibia and/or structures on the metabasitarsus, resembling the metatibial extension of *Pouillonhybos* and *Syneproctus* (Bickel, 1985: figs 7-8; Zhu et al., 2005; Runyon, 2008; Grichanov & Brooks, 2017). It is clear that not only various leg modifications have evolved convergently among empidoid lineages for subduing and handling prey, but such traits also appeared in the early phases of their diversification as evidence by their presence in at least one Cretaceous ocydromiine. The special shape of the



legs of *Pouillonhybos* suggests that it was catching its preys during flight, as for the extant Ocydromiinae (Daugeron, 1997).

Many Empidoidea are well-known for their complex mating behavior, with gifts to the female and semaphore behaviors. Nevertheless, after Downes (1969: 290), both sexes of ‘the extant Hybotinae and some Ocydromiinae’ .... ‘capture prey and the feeding habits have no relationship to the flight in the mating swarm’. Even if the extant Ocydromiinae are making swarms without mating (Daugeron, 1997), in absence of information on the morphology of the female of *Pouillonhybos*, it is not possible to infer hypotheses about the mating behavior in this ancient Hybotidae.

## Acknowledgments

We are sincerely grateful to two anonymous referees for their useful comments on the first version of the paper.

## References

- Ale-Rocha, R., 2000. Description of *Smithybos equatoriensis*, new genus and species of Hybotinae from Ecuador (Diptera, Empididae). *Revista Brasileira de Entomologia* 44, 39–42.
- Ale-Rocha, R., 2002. Revisão dos gênero *Euhybus* Coquillett (Diptera, Empididae, Hybotinae) da regio neotropical. Grupo *dimidiatus*. *Acta Amazonica* 32, 299–324.
- Ale-Rocha, R., 2007. Redescription of *Oropezella abdominalis* Collin from Chile with description of the male and a key to the Neotropical genera of Ocydromiinae (Diptera, Hybotidae). *Revista Brasileira de Entomologia* 51, 410–412.
- Ale-Rocha, R., 2008. Revision of the Neotropical genus *Lactistomyia* Melander (Diptera, Hybotidae, Hybotinae). *Zootaxa*, 1692, 1–25.



- 324 Ale-Rocha, R., De Carvalho, C.J.B., 2003. *Neohybos* gen. nov. (Diptera, Empidoidea,  
325 Hybotinae) from the Neotropical Region. *Zootaxa*, 387, 1–16.
- 326 Bezzi, M., 1904. Empididi Indo-Australiani raccolti dal Signor L. Biro. *Annales Historico-*  
327 *Naturales Musei Nationalis Hungarici* 2, 320–361.
- 328 Bezzi, M., 1909. Beiträge zur Kenntnis der südamerikanischen Dipterenfauna auf Grund der  
329 Sammelergebnisse einer Reise in Chile, Peru und Bolivia, ausgeführt in den Jahren  
330 1902–1904 von W. Schnuse. *Fam. Empididae. Nova Acta Academiae Caesareae*  
331 *Leopoldino-Carolinae Germanicae Naturae Curiosorum* 91, 293–407.
- 332 Bickel, D.J., 1985. A revision of the Nearctic *Medetera* (Diptera: Dolichopodidae). United  
333 States Department of Agriculture, Technical Bulletin 1692, 1–109.
- 334 Bigot, J.M.F., 1859. Essai d'une classification générale et synoptique de l'ordre des insectes  
335 dipteres. (7ème mémoire). Tribus des Rhabdidi et Dolichopodi (Mihi). VIIe. *Annales*  
336 *de la Société Entomologique de France* (3), 7, 201–231.
- 337 Chvála, M., 1976. Swarming, mating and feeding habits in Empididae (Diptera), and their  
338 significance in evolution of the family. *Acta Entomologica Bohemoslovaca* 73, 363–  
339 366.
- 340 Chvála, M., 1983. The Empidoidea (Diptera) of Fennoscandia and Denmark. 2. General part.  
341 The families Hybotidae, Atelestidae and Microphoridae. *Fauna Entomologica*  
342 *Scandinavica* 12, 1–279.
- 343 Cockerell, T.D.A., 1917. Insects in Burmese amber. *Annals of the Entomological Society of*  
344 *America* 10, 323–329.
- 345 Collin, J.E., 1926. Notes on the Empididae (Diptera) with additions and corrections to the  
346 British list. *Entomologist's Monthly Magazine* 62, 213–217.
- 347 Collin, J.E., 1928. New Zealand Empididae based on material in the British Museum (Natural  
348 History). London British Museum (Natural History): viii + 110 pp.

- 349 Collin, J.E. 1933. Empididae. Diptera of Patagonia and South Chile based mainly on material  
350 in the British Museum (Natural History) 4, i–viii + 1–334.
- 351 Coquillett, D.W., 1895. Revision of the North American Empidae – a family of two-winged  
352 insects. Proceedings of the United States National Museum 18, 387–440.
- 353 Cumming, J.M., Wood, D.M., 2009. Chapter 2. Adult morphology and terminology. In:  
354 Brown, B.V., Borkent, A., Cumming, J.M., Wood, D.M., Woodley, N.E. & Zumbado,  
355 M.A. (Eds.), Manual of Central American Diptera. Vol. 1. NRC Research Press,  
356 Ottawa, 9–50.
- 357 Daugeron, C., 1997. Evolution of feeding and mating behaviors in the Empidoidea (Diptera:  
358 Eremoneura). pp. 163–182. In: Grandcolas, P. (ed.). The origin of biodiversity in  
359 insects: tests of evolutionary scenarios. Mémoires du Muséum National d'Histoire  
360 Naturelle, Zoologie, Paris, 173, 354 pp.
- 361 Downes, J.A., 1969. The swarming and mating flight of Diptera. Annual Review of  
362 Entomology 14, 271–298.
- 363 Evenhuis, N.L., 1994. Catalogue of the fossil flies of the World (Insecta: Diptera). Backhuys  
364 Publishers, Leiden, 600 pp.
- 365 Grichanov, I.Ya., Brooks, S.E., 2017. 56. Dolichopodidae (long-legged dance flies). pp.  
366 1265–1320. In: Kirk-Spriggs, A.H., Sinclair, B.J. (eds). Manual of Afrotropical  
367 Diptera, Volume 2. Nematocerous Diptera and Lower Brachycera. Suricata 5. SANBI  
368 Graphics & Editing, Pretoria.
- 369 Grimaldi, D.A., Cumming, J., 1999. Brachyceran Diptera in Cretaceous ambers and Mesozoic  
370 diversification of the Eremoneura. Bulletin of the American Museum of Natural  
371 History 239, 1–124.

- 372 Grootaert, P., 1996. Species radiation of predatory empidoids (Insecta, Diptera) in lowland  
373 rainforest in northern Papua New Guinea. In: Edwards, D.S. et al. (eds.). Tropical  
374 rainforest research - Current Issues, Kluwer Academic Publishers, 183–192.
- 375 Grootaert, P., Yang, D., 2009. A new *Syndyas* Loew, 1857 (Diptera: Hybotidae: Hybotinae)  
376 from mangroves in Singapore, with a review of the Oriental and Australasian species.  
377 Raffles Bulletin of Zoology 57, 17–24.
- 378 Kato, A., 1971. On the genus *Bicellaria* Macquart from Japan (Diptera, Empididae). Kontyû  
379 39, 279–284.
- 380 Kertész, K.C., 1899. Verzeichnis einiger von L. Bíró in Neu-Guinea und am Malayischen  
381 Archipel gesammelten Dipteren. Természettudományi Füzetek 22, 173–195.
- 382 Kovalev, V.G., 1974. A new genus of the family Empididae (Diptera) and its phylogenetic  
383 relationships. Paleontological Journal 8, 196–20.
- 384 Latreille, P.A., 1809. Genera crustaceorum et insectorum secundum ordinem naturalem in  
385 familias disposita, iconibus exemplisque plurimis explicata [Tomus Quartus et  
386 Ultimas]. Koenig. Paris, France, 399 pp.
- 387 Linnaeus, C. von, 1758. Systema Naturae per regna tria naturae secundum classes, ordines,  
388 genera, species cum characteribus, differentiis, synonymis, locis. Ed. decima  
389 reformata. Holmiae, Laur. Salvii, 1, 1–823.
- 390 Liu, X.-y., Zhang, L.-l., Yang, D., 2012. Two new species of *Syneches* belonging to *S.*  
391 *signatus* species-group from Vietnam (Diptera: Empidoidea, Hybotinae). Zootaxa  
392 3300, 55–61.
- 393 Liu, X.-y., Wang, M.-q., Yang, D., 2014. Genus *Euhybus* (Diptera: Empididae) newly found  
394 in Shaanxi province with description of a new species. Florida Entomologist 97, 1598–  
395 1601.

- 396 Loew, H., 1857. Bidrag till kännedomen om Afrikas Diptera. Öfversigt at Kongliga  
397 Vetenskaps-Akademiens Förhandlingar 14, 337–383.
- 398 Loew, H., 1858. Bidrag till kännedomen om Afrikas Diptera. Öfversigt at Kongliga  
399 Vetenskaps-Akademiens Förhandlingar 15, 335–341.
- 400 Macquart, J., 1823. Monographie des Insectes Diptères de la famille des Empides, observés  
401 dans le Nord-Ouest de la France. Recueil des travaux de la Société d'Amateurs des  
402 Sciences, de l'Agriculture et des Arts à Lille 1, 137–165.
- 403 Macquart, J., 1827. Insectes Diptères du nord de la France. Platypézines, Dolichopodes,  
404 Empides, Hybotides. Lille, Imprimerie L. Danel, 1–159.
- 405 Martins-Neto, R.G., Vieira, F.M.R., Kucera-Santos, J.C., Fragoso, L.C.M., 1992. Dípteros  
406 (Insecta, Empidoidea) da Formação Tremembé, Bacia de Taubaté, Oligoceno do  
407 Estado de São Paulo. 1. Família Hybotidae. Acta Geologica Leopoldensia 15, 31–48.
- 408 Meigen, J.W., 1803. Versuch einer neuen Gattungseinteilung [sic!] der europäischen  
409 zweiflügen Insekten. Magazin für Insektenkunde, 2, 259–281.
- 410 Meigen, J.W., 1820. Systematische Beschreibung der bekannten europäischen zweiflügeligen  
411 Insekten. Forstmann, F.W. (publish.), Aachen, 2, i–x + 1–365.
- 412 Meigen, J.W., 1822. Systematische Beschreibung der bekannten europäischen zweiflügeligen  
413 Insekten. Forstmann, F.W. (publish.), Aachen, 3, i–x + 1–416.
- 414 Meigen, J.W., 1830. Systematische Beschreibung der bekannten europäischen zweiflügeligen  
415 Insekten. Forstmann, F.W. (publish.), Aachen, 6, i–iv + 1–401.
- 416 Melander, A.L., 1902. A monograph of the North American Empididae. Part 1. Transactions  
417 of the American Entomological Society 28, 195–367.
- 418 Melander, A.L., 1928. Diptera fam. Empididae. Genera Insectorum 185, 1–434.
- 419 Meunier, F., 1908. Monographie des Empidae de l'ambre de la Baltique. Annales de Science  
420 Naturelle, Zoologie, Paris (9), 7, 81–135.

- 421 Oldroyd, H. 1948. A wingless empid (Diptera) from Tasmania. Entomologist's Monthly  
422 Magazine 84, 278–279.
- 423 Papp, L., Földvári, M., 2001. A new genus and three new species of Hybotidae with new  
424 records of the Hungarian Empidoidea (Diptera). Acta Zoologica Academiae  
425 Scientiarum Hungaricae 47, 349–361.
- 426 Philippi, R.A., 1865. Aufzählung der Chilenischen Dipteren. Verhandlungen der Zoologisch-  
427 Botanischen Gesellschaft in Wien 15, 595–782.
- 428 Plant, A.R., 1989. A revision of the Ocydromiinae (Diptera: Empidoidea: Hybotidae) of New  
429 Zealand with descriptions of new genera and species. New Zealand Journal of Zoology  
430 16, 231–241.
- 431 Rafael, J.A., 1995. Revisão das espécies neotropicais de Empididae (Diptera) descritas por  
432 Mario Bezzi. III. Ocydromiinae. Revista Brasileira de Entomologia 39, 755–770.
- 433 Rafael, J.A., Cumming, J.M., 2004. The Neotropical genera *Macrostomus* Wiedemann and  
434 *Porphyrochroa* Melander (Diptera, Empididae, Empidinae). Revista Brasileira de  
435 Zoologia 21, 439–448.
- 436 Runyon, J.B., 2008. New species of long-legged flies (Diptera: Dolichopodidae) from Central  
437 Pennsylvania. Proceedings of the Entomological Society of Washington 110, 363–373.
- 438 Saigusa, T., 1986. New genera of Empididae (Diptera) from Eastern Asia. Sieboldia 5, 97–  
439 118.
- 440 Schiner, J.R., 1862. [Heft 8 (part)]. In: Fauna Austriaca. Die Fliegen (Diptera). Theil 1. Carl  
441 Gerolds Sohn, Wien, i–lxxx + 657–674.
- 442 Schlüter, T., 1978. Zur Systematik und Palökologie harzkonservierter Arthropoda einer  
443 Taphozönose aus dem Cenomanium von NW-Frankreich. Berliner  
444 Geowissenschaftliche Abhandlungen, Reihe A, Geologie und Paläontologie 9, 1–150.

- Shamshev, I.V., Wahlberg, E., Soltész, Z., 2017. New data on the genera *Allanthalia* Melander, *Chvalaea* Papp et Földvári and *Leptodromiella* Tuomikoski (Diptera: Hybotidae) from the Palaearctic. *Russian Entomological Journal* 26, 161–168.
- Shi, G., Grimaldi, D.A., Harlow, G.E., Wang, J., Yang, M., Lei, W., Li, Q., Li, X., 2012. Age constraint on Burmese amber based on U–Pb dating of zircons. *Cretaceous Research* 37, 155–163. doi.org/10.1016/j.cretres.2012.03.014
- Sinclair, B.J., 1996. Revision of the Southern Africa genus *Acarterus* Loew (Diptera: Empidoidea; Hybotinae), with description of seven new species. *Annals of the Natal Museum* 37, 215–238.
- Sinclair, B.J., 2011. Revision of Fijian *Syneches* (Diptera: Empidoidea: Hybotidae), with a reassessment of the genus. *Canadian Entomologist* 143, 358–369.
- Sinclair, B.J., 2019. Revision of the southern African genus *Stuckenbergomyia* Smith, 1971 (Diptera, Empidoidea) and proposal of a new subfamily. *African Invertebrates* 60, 133–145.
- Sinclair, B.J., Cumming, J.M., 2000. Revision of the genus *Apterodromia* Oldroyd (Diptera: Empidoidea), with a redefinition of the tribe Ocydromiini. *Records of the Australian Museum* 52, 161–186.
- Sinclair, B.J., Cumming, J.M., 2006. The morphology, higher-level phylogeny and classification of the Empidoidea (Diptera). *Zootaxa* 1180, 1–172.
- Sinclair, B.J., Cumming, J.M., 2007. *Leptopezella*, a new Southern Hemisphere genus of Ocydromiinae (Diptera: Empidoidea: Hybotidae). *Zootaxa* 1629, 27–37.
- Sinclair, B.J., Cumming, J.M., 2017. Chapter 52. Hybotidae (hybotid dance flies). In: Kirk-Spriggs, A.H., Sinclair, B.J. (eds). *Manual of Afrotropical Diptera. Volume 2. Nematocerous Diptera and Brachycera*. Suricata Pretoria, 5, 1237–1249.

- 469 Smith, K.G.V., 1961. On the genus *Macrostomus* Wiedemann (Diptera: Empididae) with  
470 notes on the included species and new synonymy. Proceedings of the Royal  
471 Entomological Society of London (B) 30, 53–56.
- 472 Smith, K.G.V., 1967. Afrikanische Empididae. (Ergebnisse der Deutschen Zoologischen  
473 Ostafrika –Expedition 1951/52, Gruppe Lindner - Stuttgart, Nr. 41) und (Ergebnisse  
474 der Forschungsreise Linder 1958/59 - Nr. 23). Stuttgarter Beiträge zur Naturkunde  
475 179, 1–16.
- 476 Smith, K.G.V., 1969. The Empididae of Southern Africa (Diptera). Annals of the Natal  
477 Museum 19, 1–347.
- 478 Solórzano-Kraemer, M.M., Delclòs, X., Peñalver, E., Sinclair, B.J., 2020. New genus and first  
479 record of Hybotinae (Diptera: Empidoidea: Hybotidae) in Middle Miocene Dominican  
480 amber. Novitates Caribaea 15, 1–8.
- 481 Steyskal, G.C., Knutson, L.V., 1981. Chapter 47 Empididae. pp. 607–624. In: McAlpine, J.F.,  
482 Peterson, B.V., Shewell, G.E., Teskey, H.J., Vockeroth, J.R. and Wood, D.M.  
483 (coordinators). Manual of Nearctic Diptera, 1, Research Branch, Agricultural Canada  
484 Monograph, Ottawa, Ontario, 27, 674 pp.
- 485 Tuomikoski, R., 1936. Bemerkungen über die Clusiiden (Dipt.) Finnlands. Annales  
486 Entomologici Fennici 2, 182–186.
- 487 Tuomikoski, R., 1959. Mitteilungen über die Empididen (Diptera) Finnlands. 6. *Trichinomyia*  
488 gen. n., eine neue Ocydromiinegattung. Annales Entomologici Fennici, 25, 103–110.
- 489 Wahlberg, E., Johanson, K.A., 2018. Molecular phylogenetics reveals novel relationships  
490 within Empidoidea (Diptera). Systematic Entomology 43, 619–636.
- 491 Walker, F., 1852. Diptera. Part III. Insecta Saundersiana: or characters of undescribed insects  
492 in the collection of William Wilson Saunders, Esq., F.R.S., F.L.S., John Van Voort,  
493 London, 1, 157–252.

- Walker, F., 1860. Catalogue of the dipterous insects collected in Amboyna by Mr. A.R. Wallace, with descriptions of new species. Journal and Proceedings of the Linnean Society of London, Zoology 5, 144–168.
- Waters, S.B., 1989. A new hybotine dipteran from the Cretaceous of Botswana. Palaeontology 32, 657–667.
- Wiedemann, C.R.W., 1817. Über einige neue Fliegen-Gattungen. Zoologisches Magazin (Wiedemann's) 1 (1), 57–61.
- Wilder, D.D. 1974. A revision of the genus *Syneches* Walker (Diptera: Empididae) for North America and the Antilles. Contributions of the American Entomological Institute 10, 1–30.
- Yang, D., Grootaert, P., 2004. A new species of *Chillcottomyia* from China (Diptera: Empididae). Transactions of the American Entomological Society 130, 165–168.
- Yin, Z.-W., Cai, C.-y., Huang, D.-y., 2018. A potentially diverse fauna of springtail-hunting scydmaenines during the late Mesozoic (Coleoptera, Staphylinidae, Scydmaeninae). Cretaceous Research 90, 163–167.
- Zhu, Y.-j., Yang, D., Masunaga, K., 2005. A review of the species of *Thambemyia* Oldroyd (Diptera: Dolichopodidae) from China. Aquatic Insects 27, 299–307.

**FIG. 1. *Pouillonhybos venator*** Ngô-Muller, Engel, & Nel, gen. et sp. nov., holotype MHNE.2020.7.1. A, habitus, ventral view. B, wing, arrow base of vein M2. Scale bars = 1 mm.

**FIG. 2. *Pouillonhybos venator*** Ngô-Muller, Engel, & Nel, gen. et sp. nov., holotype MHNE.2020.7.1. Hind leg, c.s. coxal seta, bt. basitarsus, f. femur, t. tibia, s. spur. Scale bar = 1 mm.



518 **FIG. 3. *Pouillonhybos venator*** Ngô-Muller, Engel, & Nel, gen. et sp. nov., holotype  
519 MHNE.2020.7.1. A, head, above. B, male terminalia, lateral view; ce cercus, epandrium,  
520 phallus. Scale bar = 0.5 mm (A), 0.1 mm (B).

521